

Patientinformation

	Losartan	Cozaar/Comp, Klomentan/Comp, Losarstad/Comp, Losartan, Losatrix, Losartan/Hydrochlorothiazide, Losartan/hydroklortiazid, Marozid
	Eprosartan	Teveten/Comp
	Valsartan	Diovan/Comp, Valsartan, Valsartan/hydrochlorothiazide, Valsartan/Hydroklortiazid Valsartore/Comp, Valtsu/Comp, Exforge, Entresto
	Irbesartan	Aprovel, CoAprovel, Ifirmasta, Ifirmacombi, Irbesartan, Irbesartan/hydrochlorothiazid, Karvea
	Kandesartan	Atacand/Plus, Candemox Comp, Candesarstad/Comp, Candesartan, Candesartan/Hydrochlorothiazid, Candexetil/Comp, Kairasec, Kandrozid,
	Telmisartan	Micardis/Plus, Lesamor, Telmisartan/Hydroklortiazid, Tolura, Tolucombi
Diabetes medicines		
	Metformin	Metformin, Comptact, Eucreas, Glucophage, Janumet, Jentaduetto, Komboglyze, Synjardy, Vokanamet, Xigduo
	Empagliflozin	Jardiance, Synjardy, Glyxambi
	Kanagliflozin	Invokana, Vokanamet
	Dapagliflozin	Forxiga, Xigduo, Qtern
	Ertugliflozin	Steglatro, Segluromet, Steglujan
Other medicines		

Painkillers in gel or spray form can be taken. Tablets and liquid medicines containing paracetamol (Alvedon, Pamol, Panodil, Paracetamol, Paracut, Pinex, Quramol, Theremin) can also be taken.

Never exceed the recommended dose!

Produced by the Elderly and Medication Therapy Group
(Terapigrupp Äldre och Läkemedel), August 2018, Version 2

If you are taking

MEDICINES

**that carry a risk of serious side effects in
conjunction with acute dehydration caused by**

- *Diarrhoea*
- *Vomiting*
- *High fever (over 38°C)*
- *Heatwave*
- *Acute illness resulting in difficulty eating or drinking*

Certain medicines that are used to treat heart conditions, high blood pressure, pain and diabetes could have serious side effects if they are taken when your body's fluid balance is disrupted.

Remember:

There is a risk of serious side effects from these medicines in conjunction with conditions that cause dehydration, such as vomiting, diarrhoea, fever, strong heat, or a disorder that renders you unable to eat or drink. The risk of serious side effects increases for elderly persons or individuals with reduced kidney function.

What action should be taken in the event of dehydration?

If you are taking any of the named medicines, you ought to discontinue treatment temporarily if you are dehydrated. If the problem is only temporary, treatment can recommence once it has ceased.

Contact your healthcare provider if you are in any doubt!

.....

If you are taking:

Painkillers		
	Diklofenak	Diclofenak/T, Dicuno, Diklofenak/T, Eeze, Eezeneo, Ignorin, Voltaren/T, Arthrotec/forte
	Piroxikam	Brexidol
	Tenoxikam	Alganex
	Meloxikam	Meloxicam
	Ibuprofen	Brufen/retard, Ibumax, Ibumetin, Ibuprofen, Ibuzin, Ifenin, Ipren, Nurofen Apelsin
	Naproxen	Alpoxen, Naprocur, Naprosyn Entero, Naproxen, Pronaxen, Vimovo
	Ketoprofen	Orudis/Retard
	Dexibuprofen	Tradil
	Dexketoprofen	Enantyum, Kettesse
	Celecoxib	Celebra, Celecoxib
	Etoricoxib	Arcoxia, Turox
	Nabumeton	Relifex
Diuretics		
	Bendroflumetiazid	Bendroflumetiazid, Salures, Centyl K/mite
	Hydroklortiazid	Esidrex, Hydrochlorothiazide, Hydroklortiazid
	Clortalidon	Hygropax
	Metolazon	Metolazon
	Furosemid	Furix/Retard, Furosemid, Impugan, Lasix Retard
	Bumetanid	Burinex
	Torasemid	Torasemid, Torem
	Spironolakton	Spironolactone, Spironolakton
	Eplerenon	Eplerenon, Inspra
	Amilorid	Amilorid
Medicines for heart conditions and high blood pressure		
	Digoxin	Digoxin, Lanoxin
	Kaptopril	Captopril
	Enalapril	Enalapril, Renitec, Enalapril Comp, Enalapril/Hydrochlorothiazide, Linatil Comp, Linatil Comp Mite, Renitec Comp
	Lisinopril	Lisinopril, Zestril, Lisinopril/Hydrochlorotiazid
	Perindopril	Coversyl Novum
	Ramipril	Ramipril, Triatec, Ramipril/Hydroklorotiazid
	Fosinopril	Fosinopril
	Kinapril	Accupro Comp