

Region Västra Götaland, HTA-centrum

Regional activity based HTA [Verksamhetsbaserad HTA]

Health Technology Assessment

HTA report 2019:104

Trabecular bone pattern assessment in dental radiographs for prediction of bone fracture risk

Johanen A, Bernhardsson S, Hagman J, Hakeberg M, Hange D, Jonasson G, Laine CM, Liljegren A, Lund H, Persson C, Svensson M, Stadig I, Wartenberg C, Sjögren P

Trabecular bone pattern assessment in dental radiographs for prediction of bone fracture risk [Utvärdering av trabekulärt benmönster i tandröntgenbilder för prediktion av benfrakturrisk]

Johanen A^{1,2*}, Bernhardsson S³, Hagman J^{1,4}, Hakeberg M^{1,4}, Hange D^{5,6}, Jonasson G^{1,6}, Laine CM⁷, Liljegren A⁸, Lund H⁹, Persson C², Stadig I⁸, Svensson M¹⁰, Wartenberg C³, Sjögren P³

¹ Department of Behavioral and Community Dentistry, Institute of Odontology, The Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden

² Clinic of Oral and Maxillofacial Radiology, Public Dental Service, Region Västra Götaland, Gothenburg, Sweden

³ HTA-centrum of Region Västra Götaland, Gothenburg, Sweden

⁴ Clinic of Oral Medicine, Public Dental Service, Region Västra Götaland, Gothenburg, Sweden

⁵ Department of Public Health and Community Medicine/Primary Health Care, The Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden

⁶ Närhälsan Research and Development Primary Health Care Södra Älvsborg, Region Västra Götaland, Borås, Sweden

⁷ Division of Endocrinology, Sahlgrenska University Hospital, and Institute of Medicine, University of Gothenburg, Gothenburg, Sweden

⁸ Medical Library, Sahlgrenska University Hospital, Gothenburg, Sweden

⁹ Department of Oral and Maxillofacial Radiology, University of Gothenburg, Gothenburg, Sweden

¹⁰ Health Metrics, The Sahlgrenska Academy at the University of Gothenburg, Gothenburg, Sweden

*Corresponding author

Published January 2019

2019:104

Suggested citation: Johanen A, Bernhardsson S, Hagman J, Hakeberg M, Hange D, Jonasson G, Laine C, Liljegren A, Lund H, Persson C, Svensson M, Stadig I, Svensson M, Wartenberg C, Sjögren P.

Trabecular bone pattern assessment in dental radiographs for prediction of fracture risk [Utvärdering av trabekulärt benmönster i tandröntgenbilder för prediktion av frakturrisk]. Göteborg: Västra Götalandsregionen, Sahlgrenska Universitetssjukhuset, HTA-centrum; 2019. Regional activity based HTA 2019:104

Table of contents

1.	Abstract.....	4
2.	Svensk sammanfattning – Swedish summary	5
3.	Summary of Findings	7
4.	Abbreviations/Acronyms.....	8
5.	Background.....	8
6.	Methods	12
7.	Results	12
8.	Ethical aspects	14
9.	Organisational aspects	15
10.	Economic aspects	16
11.	Discussion.....	19
12.	Future perspectives	21
13.	Participants in the project	21

Appendix 1 Study selection, search strategies and references

Appendix 2 Included studies – design and patient characteristics

Appendix 3 Excluded articles

Appendix 4 Outcome tables

1. Abstract

Background

Osteoporosis is an asymptomatic change of bone mineral density (BMD) that implies an increased risk for bone fractures. Currently, osteoporosis is an underdiagnosed and undertreated condition. Bone mineral density (BMD) measurement using dual-energy x-ray absorptiometry (DXA). DXA is not recommended for general screening due to high costs and low sensitivity. A Fracture Risk Assessment Tool (FRAX) that estimates the 10-year fracture risk is freely available online, and The Swedish National Board of Health and Welfare recommends thresholds of a FRAX-determined fracture risk of >15% for medical and radiological examination, and a FRAX-determined fracture risk of >30% for medication with bone-active drugs which can decrease the fracture risk by 40%.

Dental radiographs have been suggested as a method for identifying individuals with high fracture risk. Dental radiographs could be used for prediction of fracture risk as they are commonly taken as part of dental care in a large proportion of the population. Devices and expertise to interpret dental radiographs are already established in dental clinics.

Objectives

1. To assess whether analysis of trabecular bone pattern in an intraoral radiograph and/or panoramic radiograph can predict fracture risk in adults without diagnosed osteoporosis, compared with fracture risk prediction with DXA or with FRAX.
2. To test sensitivity and specificity for identification of osteoporosis, as compared with DXA.

Methods

A systematic literature search was conducted in April 2018 in Medline, Embase, the Cochrane Library, and several HTA-databases. The included studies were critically appraised using the QUADAS-2 tool for quality assessment of diagnostic accuracy studies and the certainties of evidence were assessed using the GRADE approach.

Main results

Two cohort studies and one cross-sectional study were included in the analysis.

The two cohort studies evaluated the use of trabecular bone pattern in dental radiographs of the lower jaw for prediction of bone fractures in women with a follow-up of 10 to 15 years. The relative risk of fracture was significantly higher for women with sparse trabecular bone pattern, identified by visual assessment of dental radiographs by dentists, in one of the studies (including two cohorts), and by digital software assessment in the other study. Visual assessment in the second study did not show significant results. Taken together, the two studies showed that trabecular bone evaluation on dental radiographs may predict fractures in adults without prior diagnosed osteoporosis. Low certainty of evidence (GRADE ⊕⊕○○).

The cross-sectional study on digital analyses of trabecular bone pattern, in relation to osteoporosis (defined as DXA BMD T-score ≤ -2.5), reported a sensitivity of 0.70 (95% CI 0.62 to 0.78), and a specificity of 0.69 (95% CI 0.64 to 0.73). Based on this study, it is uncertain whether digital image analysis of trabecular bone correlates to DXA BMD T-score ≤ -2.5 . Very low certainty of evidence (GRADE ⊕○○○).

Concluding remarks

Based on low certainty of evidence (GRADE ⊕⊕○○) trabecular bone evaluation on dental radiographs may provide a technique to predict fracture in adults without prior diagnosed osteoporosis. It is uncertain whether digital image analysis of trabecular bone correlates to DXA BMD T-score ≤ -2.5 . Very low certainty of evidence (GRADE ⊕○○○). The potential benefits of fracture risk assessments based on dental radiographs must be weighed against an increase in time consumed for assessment of dental radiographs, a growing number of referrals to DXA, as well as false positive and negative test results. In addition, information on the prevalence, test sensitivity and specificity for fracture prediction is pivotal regarding the economic consequences of the technology.

2. Svensk sammanfattning – Swedish summary

Bakgrund

Osteoporos (benskörhet) är en asymtomatisk förändring av skelettets mineralinnehåll, eller bentäthet (eng. *bone mineral density*, BMD), som innebär en ökad risk för frakturer. Idag är osteoporos ett underdiagnostiserat och underbehandlat tillstånd. Mätning av bentäthet med *Dual-energy X-ray Absorptiometry* (DXA) rekommenderas inte som screening på grund av höga kostnader och låg sensitivitet. I stället används ett frakturbedömnings-instrument, "*Fracture Risk Assessment Tool*" (FRAX), som uppskattar den 10-åriga frakturrisken och som är tillgängligt kostnadsfritt på Internet. Socialstyrelsen rekommenderar ett tröskelvärde på FRAX på > 15 % för att göra en medicinsk undersökning och på > 30 % för att sätta in behandling med benaktiva läkemedel. Frakturprebyggande behandlingar kan minska frakturrisken med upp till 40 % hos patienter som har diagnostiserats med osteoporos.

Tandröntgen har föreslagits som en metod för att identifiera individer med hög frakturrisik. Tandröntgen skulle kunna användas för att prediktera frakturrisik eftersom det är en del av sedvanlig tandvård för en stor andel av befolkningen. Både röntgenutrustning och kompetens för bedömning av röntgenbilderna finns på tandvårdsmottagningar.

Syfte

Att utvärdera huruvida analys av trabekulärt benmönster i underkäken i intraorala röntgenbilder/panoramaröntgenbilder kan användas för bedömning av frakturrisik hos vuxna utan osteoporosdiagnos, jämfört med riskbedömning med DXA eller FRAX, samt att utvärdera tandröntgenanalysernas sensitivitet och specificitet avseende osteoporos, jämfört med DXA.

Metod

Systematiska litteratursökningar utfördes i april 2018 i Medline, Embase, Cochrane Library, och flera HTA-databaser. De inkluderade studierna granskades med QUADAS-2-instrumentet för kvalitetsgranskning av diagnostiska studier och tillförlitligheten till det vetenskapliga underlaget värderades enligt GRADE.

Resultat

Två kohortstudier och en tvärsnittsstudie inkluderades i analysen.

Båda kohortstudierna utvärderade analys av trabekulärt benmönster i underkäken i tandröntgenbilder för frakturprediktion hos kvinnor (10–15 års uppföljning). Den relativa frakturrisken var signifikant högre hos kvinnor med glest trabekulärt benmönster, som identifierats genom visuell analys av röntgenbilderna av tandläkare (i den ena studien med två kohorter), och med hjälp av datorprogram i den andra studien. I den andra studien nådde inte den visuella analysen av benmönstret signifikanta resultat avseende frakturrisik. Tillsammans visar de två kohortstudierna att analys av trabekulärt benmönster i tandröntgenbilder möjligen kan användas för frakturprediktion hos vuxna utan tidigare osteoporosdiagnos. Begränsat vetenskapligt underlag (GRADE ⊕⊕○○).

Tvärsnittsstudien med digital analys av trabekulärt benmönster i tandröntgenbilder, i förhållande till osteoporos (definierad som DXA bentäthet T-värde ≤ -2.5), uppvisade en sensitivitet på 0.70 (95 % konfidensintervall 0.62 till 0.78), och specificitet på 0.69 (95 % konfidensintervall 0.64 till 0.73). Baserat på denna studie är det osäkert huruvida digital analys av trabekulärt benmönster i tandröntgenbilder korrelerar med DXA bentäthet T-värde ≤ -2.5 . Otillräckligt vetenskapligt underlag (GRADE ⊕○○○).

Sammanfattande slutsats

Begränsat vetenskapligt underlag (GRADE ⊕⊕○○) indikerar att analys av trabekulärt benmönster i tandröntgenbilder kan vara en teknik för frakturprediktion hos vuxna utan tidigare diagnosticerad osteoporos. Otillräckligt vetenskapligt underlag (GRADE ⊕○○○) indikerar att det är osäkert huruvida digital analys av trabekulärt benmönster i tandröntgenbilder korrelerar med DXA bentäthet T-värde ≤ -2.5 . Det bör beaktas att analys av tandröntgen för prediktion av frakturrisik skulle innebära en viss tidsåtgång för bedömning av röntgenbilder för tandläkare, ett ökande antal remisser till DXA, samt falskt positiva och negativa resultat som måste tas i beräkning och balanseras mot den potentiella nyttan. Prevalens, samt sensitivitet och specificitet avseende frakturprediktion är avgörande avseende de ekonomiska konsekvenserna av teknologin.

The above summaries were written by representatives from the HTA-centrum. The HTA report was approved by the Regional board for quality assurance of activity-based HTA. The abstract is a concise summary of the results of the systematic review. The Swedish summary is a brief summary of the systematic review intended for decision makers and is ended with a concluding summary.

Christina Bergh, Professor, MD

Head of HTA-centrum of Region Västra Götaland, Sweden, 19 December 2018

Quality assurance group	
Bergenheim, Anna	PT, PhD
Bergh, Christina	MD, Professor
Bernhardsson, Susanne	PT, PhD
Hakeberg, Magnus	DDS, Professor
Hansson-Olofsson, Elisabeth	RN, PhD, Senior lecturer
Jivegård, Lennart	MD, Senior university lecturer
Larsson, Anders	MD, PhD
Nelzén, Olle	MD, Associate professor
Petzold, Max	Statistician, professor
Rylander, Christian	MD, Associate professor
Sjögren, Petteri	DDS, PhD
Sjövall, Henrik	MD, Professor
Skogby, Maria	RN, PhD
Strandell, Annika	MD, Associate professor
Svanberg, Therese	HTA librarian
Svensson, Mikael	Health Economist, Professor
Wallerstedt, Susanna	MD, Professor
Wartenberg, Constanze	Psychologist, PhD

DDS	Doctor of dental surgery
MD	Medical doctor
PhD	Doctor of Philosophy
PT	Physiotherapist
RN	Registered nurse

3. Summary of Findings

Outcomes	Study design Number of studies	Relative risk (95%CI)		Certainty of evidence GRADE*
		Index test	Reference test	
Fracture prediction	1 cohort study with 15-year follow-up (n=136)	<u>Visual evaluation of sparse trabeculation</u> RR 1.52 95% CI: 0.56 to 4.11 <u>Software evaluation of bone texture</u> RR 4.74 95% CI: 1.49 to 15.04	<u>FRAX ≥7% (without BMD)</u> RR 1.34 95% CI: 0.56-3.20	⊕⊕○○ ¹
	1 cohort study with 10-year follow-up, two overlapping groups (n=499 and 412, respectively)	<u>Visual evaluation of sparse trabeculation</u> RR 2.09 95% CI: 1.3 to 3.5 RR 3.7 95% CI: 2.2-6.4	<u>FRAX >15% (without BMD)</u> RR 1.85 95% CI: 0.7 to 5.6 RR 4.1 95% CI: 2.4-7.2	
Sensitivity and specificity for DXA BMD T-score ≤ -2.5	1 cross-sectional study (n=671)	Sensitivity: 0.70 95% CI: 0.62 to 0.78	Specificity: 0.69 95% CI: 0.64 to 0.73	⊕○○○ ²

BMD = bone mineral density, CI = confidence interval, DXA = Dual-energy X-ray absorptiometry, FRAX = fracture risk assessment tool, Jaw-X: a computer software, based on an image processing algorithm, OR = Odds ratio, RR = relative risk.

Footnotes:

¹Some study limitations (patient flow not clearly described), uncertain precision (few events), and large effect (up-grade one step).

²Serious study limitations (missing data, blinding not reported, x-ray location of analysis on preference), and serious imprecision (hypothesis generating design, low prevalence).

*** Certainty of evidence**

High certainty We are very confident that the true effect lies close to that of the estimate of the effect.

⊕⊕⊕⊕

Moderate certainty We are moderately confident in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.

⊕⊕⊕○

Low certainty Confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect.

⊕⊕○○

Very low certainty We have very little confidence in the effect estimate:
The true effect is likely to be substantially different from the estimate of effect.

⊕○○○

4. Abbreviations/Acronyms

AUC	Area under curve
AR	Absolute risk
BMD	Bone mineral density
DXA	Dual-energy X-ray absorptiometry
FRAX	Fracture risk assessment tool
HTA	Health technology assessment
OR	Odds ratio
RCT	Randomized controlled trial
RR	Relative risk
ROC	Receiver operating characteristic
VGR	Region Västra Götaland
WHO	World Health Organization

5. Background

Disease/disorder of interest and its degree of severity

People with osteoporosis are at increased risk of fractures upon mechanical forces that would not ordinarily result in fracture (i.e. fragility fractures or low-trauma fractures), which may occur upon low energy injuries such as a fall (Kanis *et al.*, 2001). Osteoporosis is asymptomatic and causes no pain. Thus, the condition may not become evident until fractures occur. Some fracture risk factors cannot be altered, such as age and heredity, whereas e.g. life-style and nutrition can be modified.

Osteoporotic fractures that occur in the hip or the vertebrae are associated with high morbidity and mortality. Approximately 20% of those who suffer from a hip fracture die within the following year (Socialstyrelsen, 2012), not because of the fracture itself but due to comorbidity and/or thrombosis as a result of inactivity. Only 30% of patients regain their former physical capacity after a hip fracture. Both hip and vertebral fractures lead to decreased health-related quality of life and an increased risk of reduced physical function or immobility. Hip fracture is associated with a 20% and vertebral fractures with up to a 35% decrease in health-related quality of life (Borgström *et al.*, 2006). Vertebral fractures cause back pain and loss of height.

Although professional organizations have made recommendations for fracture prevention in different populations, there is no consensus on a gold standard for screening method. However, the osteoporosis diagnosis is commonly determined by measuring bone mineral density (BMD) with dual-energy x-ray absorptiometry (DXA) (Tu *et al.*, 2018).

BMD measured with DXA is not recommended for screening of the population because of its high costs and low sensitivity. The sensitivity for prediction of fractures varies depending on the age of the individuals from 20-70%, with the highest sensitivity for individuals over the age of 80 years (Sanders *et al.*, 2006).

The risk of fracture is highest among women with osteoporosis, but 50-70% of all fragility fractures occur in individuals with osteopenia (Pasco *et al.*, 2006; Sanders *et al.*, 2006). It is only in the very old (>80 years) that most fractures occur in women with osteoporosis. Of the women sustaining fractures, 80% of 50 to 59-year olds did not have osteoporosis, 50% of 60 to 79-year olds did not have osteoporosis, and even among those over 80 years of age, 30% did not have osteoporosis based on DXA measurements. Therefore, reducing the population burden of fractures requires attention not only to osteoporotic women but also to women with osteopenia (Sanders *et al.*, 2006).

Using the term osteoporotic fractures for fragility fractures is misleading because it implies that the fractures occur in a group of women identifiable by measurement of BMD.

The Fracture Risk Assessment Tool (FRAX) estimates the 10-year fracture risk and is freely accessible on the Internet (FRAX® Fracture Risk Assessment Tool, 2008). FRAX was developed in 2008, by Centre for metabolic bone diseases, University of Sheffield, UK. FRAX is based on clinical variables such as age, weight, previous fracture, heredity, current smoking, and use of oral glucocorticoids, rheumatoid arthritis, and alcohol intake (≥ 3 units daily). A value for BMD can be included in FRAX, but the tool can also be used without BMD. If a response is missing, the tool calculates the probability of future fracture as if the response was “no”. Inclusion of BMD increases the predictive value of FRAX. The Swedish National Board of Health and Welfare recommends a FRAX (without DXA) determined fracture risk of $>15\%$ as a threshold for medical examination and referral for BMD with DXA, and a FRAX-determined risk of $>30\%$ as threshold for medication with bone-active drugs (Socialstyrelsen, 2012).

The sensitivity and specificity levels of FRAX vary widely depending on, for example, defined population thresholds, as described in Kanis *et al.* (2007). In four studies included in a systematic review by National Clinical Guideline Centre UK (2012), FRAX (without BMD) at 10% threshold, had sensitivity ranging from 50 to 100 % (95% CI: 46% to 100%) and specificity from 0 to 72 % (95% CI: 0% to 73%), whereas at the 20% threshold the sensitivity and specificity were from 16 to 29% (95% CI: 13%–31 %), and from 81 to 93% (95% CI: 80% to 94%), respectively. At the FRAX 30 % threshold the sensitivity ranged from 4 to 10% (95% CI: 3% to 11%), and the specificity from 96 to 99% (95% CI: 95%–99%) (Bolland *et al.*, 2011; Ensrud *et al.*, 2009; Fraser *et al.*, 2011; Leslie *et al.*, 2010).

Prevalence and incidence

In 2012, the number of individuals in Region Västra Götaland who suffered an osteoporotic fracture was 5,061, and the number of individuals with a hip fracture was 1,845 (Västra Götalandsregionen, 2013). In 2017, the number of individuals in Region Västra Götaland who suffered an osteoporotic fracture was 4,784 (Västra Götalandsregionen, Vårddatabas Vega, 2019).

Present assessment

For individuals with a FRAX score of $>15\%$ (without DXA), BMD with central DXA of the hip and the lumbar vertebrae should be measured.

The normal pathway through the healthcare system and current wait time for medical assessment/treatment

The normal pathway through the healthcare system in Region Västra Götaland follows a ‘fracture liaison service’ (*frakturkedja*) in which individuals older than 50 years with an osteoporotic fracture are identified by a hospital ‘fracture chain coordinator’ and receive a treatment plan from the Osteoporosis Unit.

After the fracture, the patient may be referred to a general practitioner. They evaluate FRAX and refer for DXA (at the osteoporosis ward) if FRAX is $>15\%$. Patients can also self-refer for DXA assessment.

Present waiting time for DXA in Region Västra Götaland is 4-6 months (Sahlgrenska Universitetssjukhuset, 2018). If treatment is recommended after DXA measurements, it is possible to join a ‘School of osteoporosis’ at Sahlgrenska University Hospital, Mölndal. The general practitioner prescribes the medication and meets the patient annually. The effect of the medication is evaluated by a new DXA at least every second year. The osteoporotic ward reports the DXA results as well as their recommended time point for the next DXA assessment to the general practitioner. Treatment duration, e.g. with alendronic acid, is 4-6 years. Thereafter the medication can be paused for 1-2 years upon which a new FRAX and possibly DXA can take place.

Number of patients per year who undergo current treatment regimen

Bone-active treatments are available and decrease the fracture risk by 40% (Socialstyrelsen, 2014). Still, only 12% of patients with fragility fractures were treated with bone-specific medication between 2005 and 2012.

With the recent introduction of fracture liaison services, this proportion is likely to have increased in the subsequent years. Approximately 14% were treated with vitamin D and calcium (18% of hip fracture patients) according to the National Board of Health and Welfare in Sweden. Thus, osteoporosis is currently considered underdiagnosed and the associated risk for fracture undertreated.

Present recommendations from medical societies or health authorities

According to the National Board of Health and Welfare (Socialstyrelsen, 2014), prevention measures and/or treatment should be applied to:

- women aged at least 59 years with previous fragility/low-trauma fracture
- men aged at least 70 years with previous fragility/low-trauma fracture
- individuals with FRAX >15% (without BMD measurements)

According to regional guidelines in Region Västra Götaland (Västra Götalandsregionen, 2018), individuals older than 50 years with a fragility fracture should be identified by a hospital fracture chain coordinator and receive a treatment plan from the Osteoporosis Unit. For high risk patients, for example old people who are hospitalized due to the first hip fracture or vertebral compression, medical treatment should be initiated already before hospital discharge.

In primary care, the general practitioners should use FRAX to identify risk patients and then DXA to measure BMD, and thereafter when indicated, provide treatment with bisphosphonate (e.g. alendronic acid or zoledronic acid) or denosumab.

Health Technology at issue: Trabecular bone pattern assessment in dental radiographs as fracture risk indicator

BMD measured with DXA and FRAX are the two methods currently used for identifying individuals with an increased risk for fractures. It is a problem that even those at high risk, who e.g. already have sustained a fracture, are not necessarily assessed. Currently, efforts are made to establish new diagnostic and treatment pathways, involving various healthcare providers.

Dental radiographs have been suggested as a method for identifying individuals with high fracture risk (Sundh *et al.*, 2017; Jonasson *et al.*, 2011). There are digital as well as visual techniques for assessing fracture risk using dental radiographs. The digital techniques assess the thickness and degree of connection of the trabeculae (Lee and White, 2005; White *et al.*, 2005a), the transition from trabeculae to inter-trabecular spaces, or the number and size of the marrow spaces between the trabeculae (Jonasson and Billhult, 2013).

The visual method uses three reference images (A-C) to categorize the trabecular bone. A) sparse trabeculation (few trabeculae, large marrow spaces and an impression of more radiolucency in the image); B) heterogeneous trabeculation (both dense and sparse trabeculation where dense bone is found more adjacent to the tooth crowns and the sparse bone is more apically situated, or difficulties sorting the trabecular pattern under dense or sparse); C) dense homogeneous trabeculation (many trabeculae are connected to each other and small but many marrow spaces in the entire radiographed area).

In this HTA-report, two different types of dental radiographic images were used for bone pattern analyses: the intraoral radiograph and panoramic radiograph. The intraoral radiograph is the standard technique used by dentists and dental hygienists during regular dental check-ups. An intraoral radiograph covers a relatively limited area of the jaw bone, typically the area surrounding two to three teeth. In contrast a panoramic radiograph is obtained by scanning a larger area covering the upper and lower jaws and surrounding tissues, basically from ear to ear, and gives an overview of a large area. However, the intraoral radiograph has a higher resolution compared with the panoramic radiograph and is therefore more detailed.

The advantages of using dental radiographs as a screening tool for the assessment of fracture risk may be:

- A large proportion of the adult population visit their dentist annually
- Individual intraoral radiographs taken on a regular basis in dental examinations use largely similar projections and exposures, making them suitable for comparisons over time.
- All dental care clinics already have intraoral X-ray devices. Panoramic radiographic equipment is however not found at a majority of general dental clinics.
- Dentists are well trained in interpreting dental radiographs and prevention is one of the dentists' primary tasks.
- Periapical radiograph is the type of image that permits evaluation of the trabecular bone well besides diagnosis of jaw infections usually from the root canal/pulp system of teeth, while bitewing radiographs images tooth crowns and cervical part of the trabecular bone, close to the tooth crown. Therefore, these radiographs are mostly used for diagnosis of caries and periodontitis. A panoramic radiograph reveals both jaws, including the mandibular cortex and all present teeth as well as other anatomic structures within the oral and maxillofacial area.
- Periapical radiographs are used regularly during dental examinations, albeit on diagnostic indications.
- The remodelling rate of the trabecular bone is much higher than that of the cortical bone (Kanis, 1994). Furthermore, the remodelling rate of the mandibular alveolar bone is generally higher than that of the trabecular bone in other parts of the skeleton (Huja *et al.* 2006), which facilitates fracture risk detection using dental radiographs.

Objective

The main question for the current HTA project

Can analysis of trabecular bone pattern in an intraoral radiograph and/or panoramic radiograph predict fracture risk in adults without diagnosed osteoporosis, compared with DXA or FRAX?

PICO: (*P=Patient I=Intervention C=Comparison O=Outcome*)

P	Adults without diagnosed osteoporosis
I	1: Analysis of trabecular pattern in an intraoral radiograph 2: Analysis of trabecular pattern in a panoramic radiograph
C	1: Dual-energy X-ray absorptiometry (DXA) value 2: Fracture risk assessment tool (FRAX) score
O	<p><u>Critical for decision making</u> Fracture prediction</p> <p><u>Important for decision making</u> Test sensitivity and specificity for osteoporosis defined as with DXA T-score ≤ -2.5</p> <p><u>Less important for decision making</u> Risks and/or complications (not related to radiation)</p>

For a detailed description, see Appendix 1.

6. Methods

Systematic literature search (Appendix 1)

During April 2018 two authors (IS, AL) performed systematic searches in Medline, Embase, the Cochrane Library and a number of HTA-databases. Reference lists of relevant articles were also scrutinised for additional studies. Search strategies, eligibility criteria and a graphic presentation of the selection process are presented in Appendix 1. These authors conducted the literature searches, selected studies, and independently of one another assessed the obtained abstracts and made a first selection of full-text articles for inclusion or exclusion. Any disagreements were resolved in consensus. The remaining articles were sent to all the participants of the project group. All authors read the articles independently of one another and it was finally decided in a consensus meeting which articles should be included in the assessment.

Critical appraisal and certainty of evidence

The included studies, their design, and patient characteristics are presented in Appendix 2. Excluded studies, with reasons for exclusion, are presented in Appendix 3. The included studies have been critically appraised using the QUADAS-2 tool for quality assessment of diagnostic accuracy studies (Whiting *et al.*, 2011; QUADAS-2, n.d.). The findings of each study have been summarized per outcome in Appendix 4. A summary result per outcome and the associated certainty of evidence are presented in a Summary-of-findings table (page 7). Certainty of evidence was assessed according to the GRADE approach (Atkins *et al.*, 2004; GRADE Working group).

Ongoing research

A search in www.clinicaltrials.gov (2018-09-05) was conducted with the keywords:

(Dental Radiography OR Intraoral OR Intra-oral OR Periapical OR Peri-apical OR Periapex OR Peri-apex OR PanORamic OR Pantomography OR Pantomographies OR Orthopantomography OR Orthopantomographies OR Panoramic Radiography OR Panoramic Radiographies OR Radiovisiograph OR Scanora OR Scanoras OR Visualix OR Visualices OR Digora OR Digoras OR Sens-A-Ray OR Sens A Ray OR SensARay)

AND

(Cancellous or Spongy or Trabecular OR Intertrabecular or Bone Density OR Bone Densities OR Bone Mineral Density OR Bone Mineral Densities OR BMD or Bone mineral content OR Bone Mineral Contents OR Bone texture).

7. Results

Systematic literature search (Appendix 1)

The literature search identified 2,377 articles after removal of duplicates. After reading the abstracts 2,301 articles were excluded. Another 34 articles were excluded by two authors after reading the articles in full text. The remaining 42 articles were sent to all participants of the project group, and three articles (two cohort studies and one cross-sectional) were finally included in the assessment (Appendix 2).

Results

Three publications, reporting on two cohort studies and one cross-sectional study, met the inclusion criteria for the central question (Jonasson and Billhult, 2013; Sundh *et al.*, 2017; Verheij *et al.*, 2009).

The two cohort studies had fracture as outcome (Jonasson and Billhult, 2013; Sundh *et al.*, 2017). Both studies had low precision mainly due to small to moderate sample sizes and few fracture events.

The cross-sectional study (Verheij *et al.*, 2009) calculated sensitivity and specificity for using a digital image analysis of trabecular bone pattern in the mandible as an indicator of BMD, which was assessed by DXA scan of lumbar spine and hip.

This study employed a hypothesis-generating design, in which several different markers were used to identify the potentially best sensitivity and specificity of osteoporosis diagnosis, thereby introducing a risk for significant results by chance. Furthermore, the regions of interest in the dental radiographs were selected manually and blinding of the assessments was not reported.

Analysis of trabecular bone pattern in intraoral or panoramic radiographs to predict fracture risk in adults compared with DXA

Outcomes, critical for decision-making

Fracture (Appendix 4.1)

Two cohort studies that compared evaluation of trabecular bone pattern in intraoral or panoramic radiographs with DXA and/or FRAX reported fracture as an outcome. Both studies had limitations regarding precision (see above). The settings differed between the studies. In the cohort study by Jonasson and Billhult (2013) the information regarding the studied population was unclear with regard to sample selection and exclusion criteria.

The cohort study by Jonasson and Billhult (2013), with up to 15-year follow-up, reported relative fracture risks assessed with different methods (correlated to subsequent fracture rates).

The computer-based assessment of trabecular bone in intraoral radiographs resulted in a RR of fracture of 4.74 (95% CI: 1.49 to 15.04, $p < 0.05$), and FRAX combined with BMD (DXA forearm) in a RR of fracture of 2.53 (95% CI: 1.13 to 5.63, $p < 0.05$). The other fracture risk assessments were not statistically significant (e.g. visual evaluation, software tool Jaw-X, and FRAX without BMD). The absolute fracture rate was 19.1% in the total study population.

The cohort study by Sundh *et al.* (2017), studied two population samples of randomly selected women over a mean period of 10 years (a substantial number of women participated in both cohorts). The fracture rates during the follow-up periods were compared between the groups that were considered at risk (FRAX > 15% and/or sparse trabeculation), with those without identified fracture risk factors.

With visual assessments of panoramic radiographs for detection of sparse trabeculation, for fracture prediction, in the first cohort (years 1980-1992, age 50-66 years, 499 women), the RR for fractures was 2.09 (95% CI: 1.3 to 3.5). In the second cohort (1992-2002, age 62-78 years, 412 women), the RR for fractures was 3.7 (95% CI: 2.2 to 6.4). For fracture prediction with FRAX alone (threshold >15%) the corresponding RRs were 1.85 (95% CI: 0.7 to 5.6) in the 1980-1992 cohort, and 4.1 (95% CI: 2.4 to 7.2) in the 1992-2002 cohort, respectively.

The absolute fracture rates were 14.2% in the 1980-1992 cohort and 16.3% in the 1992-2002 cohort.

Conclusion

Trabecular bone evaluation on dental radiographs may be used to predict fracture in adults without a prior diagnosis of osteoporosis. Low certainty of evidence (GRADE ⊕⊕○○)

Outcomes, important for decision-making

Test sensitivity and specificity for osteoporosis measured with DXA (Appendix 4.2)

One cross sectional study (Verheij *et al.*, 2009) reported sensitivity and specificity of trabecular bone pattern assessment in intraoral- and panoramic radiographs (as index test) for prediction of osteoporosis, defined as DXA obtained BMD T-score $\leq - 2.5$.

The study employed a hypothesis-generating design and had several study limitations (see above), and the mean age of the study population was 54.8 years (SD 6.1).

Regarding morphological, digital analyses of trabecular bone pattern, in relation to DXA BMD T-score ≤ -2.5 , the sensitivity for osteoporosis, defined as BMD T-score ≤ -2.5 , was 0.70 (95% CI: 0.62 to 0.78), and the corresponding specificity was 0.69 (95% CI: 0.64 to 0.73).

Conclusion

It is uncertain whether digital image analysis of trabecular bone correlates to DXA BMD T-score ≤ -2.5 . Very low certainty of evidence (GRADE $\oplus\text{OOO}$).

Outcomes, less important for decision making

Risks and/or complications (not related to radiation)

No studies reported complications.

8. Ethical aspects

For patients diagnosed with osteoporosis, currently available treatments have the potential to improve health-related quality of life and longevity. However, implementing the proposed new test method, with analyses of intraoral radiographs, will lead to increased referrals from dental care to primary care, more healthcare visits and laboratory tests. This could affect the availability of primary care for other medical conditions. For dental health care providers, implementing the new method would add another diagnostic task to the ordinary dental examination which may lead to inappropriate prioritisations.

Due to the potential harmful effects of ionizing radiation, every radiographic examination must be justified (one extra intraoral radiograph with effective dose of $<1.5\mu\text{Sv}$ (Ludlow, 2008), is comparable to 1 day of natural background radiation). If one assumes that assessment of trabeculation is performed on already existing radiographs, the proposed method will not add any extra exposure, and it would be valuable to be able to utilize the information obtained from dental radiographs for an additional purpose, such as prediction of fracture risk.

What are the risks, from an individual perspective, given a false positive assessment of osteoporosis? This question is highly dependent on how these individuals will be taken care of in the public health care system. Would it add further, unnecessary, laboratory tests and/or examinations? Regarding radiation risk, the exposure to radiation from a BMD examination using DXA leads to an effective radiation dose in the range of $2.4\text{--}13.3\mu\text{Sv}$ (i.e. $0.0024\text{--}0.013\text{ mSv}$), depending on equipment, region and tissue weighting factors used (Blake *et al.*, 2006). This radiation dose is minuscule in relation to the annual mean radiation dose in Sweden of 2.4 mSv (for non-smokers) (Andersson, 2007). The radiation dose from one intraoral dental radiograph is about 0.01 mSv (Andersson, 2007).

In Sweden DXA examinations are conducted based on medical history and clinical examination and thus, only individuals at risk for disease are examined. Another issue to be considered is how information given by a dentist could affect the psychological well-being of an individual, who came for an ordinary dental examination, but got unexpected information about a hypothetical diagnosis of osteoporosis and then must wait several months for a final assessment regarding this diagnosis.

The benefit of the new prediction method for fractures (or osteoporosis) in dental care may be clear given the high number of older individuals/patients attending dental care on a regular basis. The method does not discriminate between gender, ethnicity or age, thus not violating ethical principles in health care, but individuals that do not attend to dental check-ups on a regular basis would be missed.

However, prediction of fractures in dental practices would imply that resources in dental care and health care are used for the new technique instead for other conditions needing attention. This possible scenario must be weighed against the advantage of fracture prediction - reducing the number of skeletal fractures in the older population and reducing the cost for treatment and suffering for individuals, but also in a societal perspective.

9. Organisational aspects

Time frame for the putative introduction of the new health technology

Currently we are mainly working with secondary fragility fracture prevention, since osteoporosis is most commonly suspected only after an individual already has sustained a fracture. With the new technology we would aim for primary prevention – i.e. to identify patients before their first fracture. Introducing a new fracture prediction method may take a long time.

Present use of the technology in other hospitals in Region Västra Götaland

Evaluation of trabecular bone pattern on dental radiographs for the purpose of assessing osteoporosis-related fracture risk is presently not used in clinical practice in Region Västra Götaland, nor elsewhere to our knowledge.

The guidelines from the National Board of Health and Welfare today recommend that when osteoporosis is suspected, the healthcare services should assess the fracture risk by FRAX and, if necessary, conduct a DXA (Socialstyrelsen, 2014).

Consequences of the new health technology for personnel

A majority (about 80-85%) of older adults make regular visits to their dental care clinics. Today dental intraoral (periapical) radiographs are already used as a complement to the examination of the teeth and the alveolar bone.

A closer communication and collaboration between the dental care clinics and the medical professions, in the first line Primary Health Care Centres, will be necessary if the new technology is introduced and when osteoporosis can be suspected from evaluation of the radiographs. If individuals with sparse bone structure on dental radiographs are referred for evaluation to Primary Health Care Centres, there will be an increase in the number of visits, DXA screenings, and blood tests to exclude secondary osteoporosis. The need for DXA should first be assessed based on medical history, symptoms and FRAX-score. Early identification of individuals with osteoporosis would enable doctors to treat causes of secondary osteoporosis or initiate bone-strengthening medication timely, thereby preventing a number of osteoporosis-related fractures. A screening visit, including clinical examination and reviewing the medical history and possible symptoms of the individual, may limit the need for DXA screenings.

Consequences for other clinics or supporting functions at the hospital or in the Region Västra Götaland

A more widespread use of dental radiographs for prediction of fractures is likely to increase the demand for resources in the Primary Health Care Centres in the short term. Also, a possible organisational challenge is the need of communicating information for several patients between dental- and primary care. However, in the long term, introduction of the new technology intends to increase the detection and treatment of osteoporosis. This would be expected to include health improvements in patients suffering from osteoporosis as well as a better health-related quality of life, together with a greater functional ability and a reduced number of fractures (Socialstyrelsen, 2014). For health care, the preventive work by dental and primary care would aim to reduce the number of fractures that need to be treated in orthopaedic surgery and rehabilitation.

10. Economic aspects

Present costs of currently used technologies

The current technologies used for identifying patients at risk for osteoporotic fracture are FRAX and BMD measurement using DXA, with costs of about 1,500 SEK per physician visit including a FRAX, and an additional 1,000 SEK per DXA (which requires the associated visit, thus a total cost of 2,500 SEK for DXA measurement).

The intervention (dental intraoral radiographs) is intended to substantially increase the number of analysed individuals, which implies that for the large mass of individuals analysed, there is no current alternative technology used and thus no associated costs. Initially, the intervention will therefore be cost-increasing compared to the current technology. At a second stage, analyses of dental radiographs for fracture prediction, earlier diagnosis and treatment could potentially reduce the costs associated with hip and other fractures.

Model of the economic consequences of the new technology

Figure 1 shows a simple model of the economic consequences of introducing dental intraoral radiographs with the associated resource use (costs) in each path of the model. In our analysis we assume a situation in which trabecular bone pattern assessment in dental radiographs would be implemented for the population aged 60 and above in Region Västra Götaland.

The first event is the additional time spent for analysis of the dental intraoral radiograph, which is estimated to consume approx. 5 minutes of dentist time per screening, and if a patient at risk is identified an additional 10 minutes of dentist time to write a referral. The dentist time cost is based on the average wage of dentists including social fees and mandatory (for the employer) retirement savings, i.e. the full market price per dentist hour. For an individual with a referral, additional costs will result due to a primary care visit, lab tests, and potentially also a DXA. If treatment is initiated, the drug treatment regimen includes Alendronate or Zoledronic acid and Calcium with Vitamin D.


Finally, an important aspect from an economic perspective is the potential savings that may materialise if currently untreated individuals with osteoporosis are identified and provided preventive treatment. Fractures, especially hip fractures, come with a substantial health care cost, and bone-active treatments have been shown to reduce the risk of hip and other fractures.

The challenge with an economic analysis of introducing dental intraoral radiographs is that for many of the relevant epidemiological and clinical variables there are great uncertainties regarding the magnitude of the effects, for example:

- What is the prevalence of currently unidentified osteoporosis patients that could potentially be diagnosed and treated based on introducing dental intraoral radiographs?
- What is the sensitivity and specificity of the dental intraoral radiographs for identification of osteoporosis?
- What is the risk reduction in terms of fewer incident fractures for patients who would receive drug treatment?

Considering the uncertainties, we present the economic consequences both as intervals including 95% of the most likely outcomes as well as the result in different scenarios.

Figure 1. Overview of cost consequences



Assumptions for the economic analysis

As described, there are many uncertainties related to the economic consequences of introducing dental intraoral radiographs, and Table 1 lists the variables that is part of the economic analysis and also the assumptions used to produce the results in this report.

Some variables are assumed to be relatively certain and have a modest impact on the result (e.g. dentist screening time of 5 minutes), whereas there are substantial uncertainties in other cases (e.g. prevalence of undiagnosed osteoporosis patients, the sensitivity and specificity of the dental intraoral radiographs, fracture risk reduction if initiating drug treatment, etc.). The variables with large uncertainty and with substantial impact on the results were analysed using a range of reasonable values, and the uncertainty of these analyses are shown using 95% credible intervals.

Cost per reduced fracture

Table 2 shows the cost per each reduced fracture if introducing dental intraoral radiographs and how it varies with different scenarios/assumptions. The mean estimate is based on the values listed in Table 1, and for variables with uncertainties it is based on the mid-point of the uncertainty range in Table 1.

The different scenarios show how the cost varies when the value of a parameter is changed. The mean estimate shows that the cost per reduced fracture is 28,300 SEK, which implies that for each reduced fracture the health care costs increase by 28,300 SEK. The reduced costs due to fewer fractures are thus not enough to compensate for the increased costs due to analysing, diagnosing and screening additional individuals.

Table 1. Assumptions for the economic analysis

Variable	Value for primary analysis
Dentists' time use costs per hour (incl. social fees)	410 SEK
Dental Intraoral Radiograph screening time	5 min
Referral time for identified individuals	10 min
Prevalence of undiagnosed osteoporosis	3% to 17%
Sensitivity for identification of osteoporosis	60% - 80%
Specificity for identification of osteoporosis	60% - 80%
Cost per physician visit	1,500 SEK
Cost of lab tests	500 SEK
Proportion of DXA for individuals with referral	50% to 100%
Cost per DXA	1,000 SEK
Proportion of identified patients (with DXA) receiving/complying treatment	60% to 100%
Alendronic acid	198 SEK/year
Zoledronic acid	2608 SEK/year
Calcium with Vitamin D	588 SEK/year
Years of drug treatment	5 years
Fracture risk reduction of drug treatment	30% - 40%
Savings per prevented fracture (adjusted for prevalence of different fractures)	55,000 SEK

Table 2. Cost per reduced fracture

Scenario	Cost per reduced fracture
Mean estimate	28 300 SEK
Prevalence of undiagnosed osteoporosis patients is (instead of 10%):	
...3%	141 900 SEK
...17%	8 200 SEK
Proportion of diagnosed patients receiving and complying with treatment is (instead of 80%):	
...60%	46 600 SEK
...100%	17 600 SEK
Fracture risk reduction of drug treatment is (instead of 35%):	
...30%	42 200 SEK
...40%	17 800 SEK
Sensitivity and specificity for identification of osteoporosis is (instead of 70%):	
...60%	52 000 SEK
...80%	10 400 SEK

The results in Table 2 for the different scenarios show that the cost per reduced fracture varies from 8,200 SEK to 141,900 SEK. The variable with the largest impact is the prevalence of undiagnosed osteoporosis. A relatively higher prevalence implies a much lower cost per reduced fracture, i.e. an economically more beneficial outcome. This is a representation of the fact that dental radiographs as a screening method is more economically motivated in populations with a higher level of undiagnosed osteoporosis. Regarding the sensitivity and specificity, the latter is the most important from an economic perspective. The reason is that false positive results (poor specificity) will create an increased volume of patients to primary care for evaluation where there is no benefit.

Cost per screened person (budget impact)

Table 3 shows the cost per screened person, which can be seen as the budget impact of introducing dental radiographs in the population aged 60 and above. The results are shown from a dental care perspective, which only considers dental care cost consequences, and a health care perspective that only considers health care cost consequences. Table 3 also shows the joint health care and dental cost consequences with and without taking future fracture risk reductions into consideration.

The results show the interval of the expected mean cost per each person screened based on the assumptions (with uncertainties) listed in Table 1.

Table 3. Budget impact per screened person with dental intraoral radiograph

Perspective	Cost per screened person (95% credible interval)
Dental care	57 SEK
Health care	1 060 SEK to 2 190 SEK
Dental and health care <u>with</u> fracture risk reduction	130 SEK to 970 SEK

From a dental care perspective, the new technology will imply a cost at around 57 SEK per person screened. Hence, if introducing dental intraoral radiographs each dental visit including a screening will be expected to cost 57 SEK more.

The largest cost impact is seen at the health care perspective where each person screened will increase costs by 1,060 to 2,190 SEK per screened person. This is mainly driven by the inflow of additional physician visits, tests and (for some) prescription drug expenses. The large interval of the expected cost increase is caused by the great uncertainties in mainly the underlying epidemiological and clinical data on the chain of consequences with dental intraoral radiographs.

Finally, the joint dental and healthcare perspective accounting for future reductions in fracture risk shows that the cost per screened person will be in the range of 130 to 970 SEK.

Possibility to adopt and use the new technology within the present budget

The results indicate that it will not be possible to fit the new technology within the present budget. Especially the increase in primary care visits due to identifying more at-risk patients and the increase in prescription drugs use will require additional budget funds, or it will displace some of the currently provided primary care.

Available economic evaluations or cost advantages/disadvantages

No economic evaluations or cost-consequence or budget-impact analysis studies of the new technology were identified in the published literature.

11. Discussion

The present HTA report identified a lack of longitudinal investigations with fracture as an outcome. Only two studies fulfilled the inclusion criteria. In addition, diagnostic studies with osteoporosis as outcome were virtually absent, and only one study fulfilled the inclusion criteria. All three studies included solely females. Based on low certainty of evidence (GRADE ⊕⊕○○), trabecular bone evaluation on dental radiographs may predict fracture in adults without prior diagnosed osteoporosis. It is uncertain whether digital image analysis of trabecular bone correlates to DXA BMD T-score ≤ -2.5 (very low certainty of evidence, GRADE ⊕○○○).

Very few fracture events were reported by Jonasson and Billhult (2013). Several participants increased their bone density (BMD measured by DXA) during the study period, by improving nutrition and physical activity after information was given. Osteoporotic participants were referred to their respective primary care centre or general practitioner for further examination and were prescribed medication calcium and vitamin D supplements and sometimes oestrogen too.

At the time of the study's baseline (1996), no national guidelines were available for the treatment of osteoporosis nor any established diagnostic criteria.

FRAX and visual trabecular pattern could be easily assessed in the dental clinic. Although beyond the scope of this report, it seems that combining assessments of mandibular bone trabeculation with other risk factors could increase the fracture predictive value substantially, especially in older individuals (Sundh *et al.*, 2017).

Having no risk factor was associated with a very low risk for fracture (1.5%), whereas the relative risk of fractures increased by 16 to 23 times when combining various risk factors (Sundh *et al.*, 2017). By comparing the gradient of risks reported by Sundh *et al.* (2017) and that reported by Kanis *et al.* (2007), it seems that the combination of a FRAX score of > 15% and the finding of mandibular sparse trabeculation in dental radiographs could predict major osteoporotic fractures to approximately the same extent as FRAX combined with BMD measurements using DXA (Kanis *et al.*, 2007; Sundh *et al.*, 2017).

Sensitivity and specificity for sparse trabeculation as fracture predictor were 27% and 88%, respectively (Jonasson and Billhult, 2013). These rates are comparable to the sensitivity and specificity for osteoporosis found in a study by Lindh *et al.* (2008) (sensitivity 28.2% and specificity 90.8%) where the trabecular pattern was evaluated as indicator of osteoporosis.

In the younger cohort (Sundh *et al.*, 2017), sparse trabeculation as fracture predictor had a sensitivity of 61% and specificity 66%. For FRAX, sensitivity was 25% and specificity 94% (without DXA). In the older cohort, sparse trabeculation as fracture predictor had a sensitivity of 78% and specificity 57%, whereas for FRAX, sensitivity was 79% and specificity 58% (Sundh *et al.*, 2017).

There is no gold standard established for fracture prediction (except the subsequent outcome itself). However, BMD measured by DXA of the hip and FRAX are currently the most widely investigated methods and were therefore chosen as reference standards (comparisons) in this report.

In order to obtain the best directness regarding the clinical setting of general dentistry, the population of interest for this report was defined as "adults without diagnosed osteoporosis". The technology at issue was "analysis of trabecular pattern in dental radiographs". Again, for best directness, as well as clinical relevance, evaluations of cortical bone were omitted because general dental practices may not have panoramic radiographic equipment. Panoramic radiographs also have lower resolution than intraoral radiographs and more practice is needed to perform trabecular assessment with panoramic radiographs than with periapical radiographs. Thus, for this purpose only periapical radiographs are suitable.

Currently used methods of analysing the trabecular bone of the mandible may not be better for detecting the risk of developing osteoporosis and risk for fragility fractures than the standard methods used in common examination pathways in general health care, such as FRAX and DXA. Thus, a knowledge gap is present as of today, however new emerging analysis tools based on digital algorithms will be developed which may further improve predictive values.

Consequence of assessment of trabecular bone pattern in dental radiographs for fracture risk are that more time would be required for this assessment, the number of referrals to primary care for medical examinations and eventually for DXA would increase, and a number of false positive and negative test results would be expected - all of which should be taken into account and balanced against the potential benefits. The prevalence, test sensitivity and specificity for fracture prediction is pivotal regarding the economic consequences of the technology.

12. Future perspectives

Scientific knowledge gaps

Larger prospective studies in varying settings with fracture as outcome are warranted. One study from the Prospective Population Study of Women in Gothenburg, Sweden (Jonasson *et al.*, 2011, 2013) with the same data set as Sundh *et al.* (2017) also demonstrates the potential benefit of using dental radiographs as fracture predictors. Long-term studies assessing trabecular pattern in intraoral and panoramic radiographs are also needed, as well as comparisons with up-to-date FRAX and DXA measures. Furthermore, existing research lacks data on men as well as on populations with a wide range of older ages. Because people today tend to live longer, we need a better understanding of the potential role of trabecular pattern assessment.

Despite advances and ongoing intense efforts to evaluate and develop optimal software algorithms for artificial intelligence, gaps remain in this area before it can be applied in dental care. Several research groups are developing varying texture analysis methods for measuring trabecular volume, spacing, and connectivity by computed tomography, magnetic resonance, and cone beam computed tomography (Geraets, Lindh and Verheij, 2012; Geraets *et al.*, 2007; Kavitha, 2015, 2016; Kathirvelue and Anburajan, 2014; Koh *et al.*, 2012; Licks *et al.*, 2010; Lee and White, 2005, Lurie *et al.*, 2012; White *et al.*, 2005a; Brasileiro *et al.*, 2017; Mostafa *et al.*, 2016), but presently the cost and complexity of these methods limit their utility for clinical routine and as a screening method.

Ongoing research

A search in www.clinicaltrials.gov (2018-09-05) yielded 48 records. None of the trials were relevant for the question at issue.

13. Participants in the project

The question was nominated by

Magnus Hakeberg, DDS, Professor, Department of Behavioural and Community Dentistry, Institute of Odontology, The Sahlgrenska Academy, University of Gothenburg and Clinic of Oral medicine, Public Dental Service, Region Västra Götaland, Gothenburg, Sweden.

Participating healthcare professionals

Astera Johanan, DDS, PhD student, Clinic of Oral and Maxillofacial radiology, Gothenburg, Public Dental Service, Region Västra Götaland and Department of behavioural and community dentistry, Institute of odontology, Sahlgrenska Academy, University of Gothenburg, Sweden.

Magnus Hakeberg, DDS, Professor, Department of Behavioural and Community Dentistry, Institute of Odontology, The Sahlgrenska Academy, University of Gothenburg and Clinic of Oral medicine, Public Dental Service, Region Västra Götaland, Gothenburg, Sweden.

Grethe Jonasson, DDS, associated professor, Närhälsan Research and Development Primary Health Care, Borås and Department of behavioural and community dentistry, Institute of Odontology, Sahlgrenska Academy, University of Gothenburg, Sweden.

Henrik Lund, DDS, PhD, Odont.Dr, Senior lecturer, Consultant Oral & Maxillofacial Radiology, Department of Oral and Maxillofacial Radiology, Institute of Odontology, Sahlgrenska Academy, University of Gothenburg, Sweden.

Jennie Hagman, DDS, PhD student, Clinic of Oral Medicine, Public Dental Service, Region Västra Götaland and Department of behavioural and community dentistry, Institute of Odontology, Sahlgrenska Academy, University of Gothenburg, Sweden.

Dominique Hange, MD, associated professor, Department of Public Health and Community Medicine/Primary Health Care, Sahlgrenska Academy, University of Gothenburg and Närhälsan Research and Development Primary Health Care, Borås, Sweden

Christine M. Laine, MD, PhD, Division of Endocrinology, Sahlgrenska University Hospital, and Institute of Medicine, University of Gothenburg, Gothenburg, Sweden

Cecilia Persson, DDS, Post graduate student. Clinic of Oral and Maxillofacial radiology, Gothenburg, Public Dental Service, Region Västra Götaland, Sweden.

Participants from the HTA-centrum

Petteri Sjögren, DDS, PhD,

Susanne Bernhardsson, Physiotherapist, PhD,

Constanze Wartenberg, Psychologist, PhD

- all at HTA-centrum, Region Västra Götaland, Gothenburg, Sweden

Ida Stadig, Librarian,

Ann Liljegren, Librarian

- both at the Medical Library, Sahlgrenska University Hospital, Region Västra Götaland, Gothenburg, Sweden.

Mikael Svensson, professor, Health Metrics, The Sahlgrenska Academy, University of Gothenburg, Sweden.

External reviewers

Göran Kjeller, Associate Professor, Department of Oral & Maxillofacial Surgery Institute of Odontology The Sahlgrenska Academy, University of Gothenburg, Sweden

Robert Eggertsen, MD, PhD, professor, Department of Medicine, University of Gothenburg, Mölnlycke Primary Health Care Centre, Mölnlycke, Sweden

Declaration of interests

Grethe Jonasson, Dominique Hange and Magnus Hakeberg are co-authors of the publication Sundh *et al.*, 2017.

Grethe Jonasson also is co-author of the publication Jonasson and Billhult, 2013.

Project time

The HTA was accomplished during the period of April to December 2018.

Literature searches were made in April 2018.

Appendix 1: Search strategy, study selection and references

Question(s) at issue:

Can analysis of trabecular bone pattern in an intraoral radiograph and/or panoramic radiograph predict fracture risk in adults without diagnosed osteoporosis, compared with DXA or FRAX?

PICO: (*P=Patient I=Intervention C=Comparison O=Outcome*)

P1	Adults without diagnosed osteoporosis
I	1: Analysis of trabecular pattern in an intraoral radiograph 2: Analysis of trabecular pattern in a panoramic radiograph
C	1: Dual-energy X-ray absorptiometry (DXA) value 2: Fracture risk assessment tool (FRAX) score
O	<u>Critical for decision making</u> Fracture prediction <u>Important for decision making</u> Test sensitivity and specificity for osteoporosis defined as with DXA T-score ≤ -2.5 <u>Less important for decision making</u> Risks and/or complications (not related to radiation)

Eligibility criteria

Study design:


Systematic reviews
Randomised controlled trials
Cohort studies
Case series ≥ 500 patients

Language:

English, Swedish, Danish, Norwegian, Finnish

Publication date: 1990-

Selection process – flow diagram



Database: Ovid MEDLINE(R) Epub Ahead of Print, In-Process & Other Non-Indexed Citations, Ovid MEDLINE(R) Daily and Ovid MEDLINE(R) 1946 to Present

Date: 2018-04-13

No. of results: 1822 ref.

#	Searches	Results
1	exp Radiography, Dental/	20222
2	((Dental or Intraoral or Intra-oral or periapical or peri-apical or Periapex or Peri-apex or tooth or teeth or mandib* or maxilla* or jaw*) and (radiograph* or radiolog* or x-ray* or xray* or imag*)).ab,ti.	51183
3	(Panoramic* or Pantomogra* or Orthopantomogra* or radiovisiograph* or Scanora* or Visualix or Visualices or Digora* or Sens-A-Ray* or Sens A Ray* or SensARay*).ab,ti.	8385
4	1 or 2 or 3	64416
5	exp Cancellous Bone/	423
6	exp Bone Density/	48104
7	(Cancellous or spongy or trabecul* or Intertrabecul* or Bone densit* or Bone mineral densit* or BMD or Bone mineral content* or bone textur*).ab,ti.	91404
8	5 or 6 or 7	103817
9	4 and 8	2488
10	(animals not (animals and humans)).sh.	4413620
11	9 not 10	1998
12	(comment or editorial or letter).pt.	1621349
13	11 not 12	1992
14	limit 13 to yr="1990 -Current"	1927
15	limit 14 to (danish or english or finnish or norwegian or swedish)	1822

Database: Embase 1974 to 2018 April 12

Date: 2018-04-13

No. of results: 1960 ref.

#	Searches	Results
1	exp dental radiology/	17537
2	exp panoramic radiography/	9114
3	((Dental or Intraoral or Intra-oral or periapical or peri-apical or Periapex or Peri-apex or tooth or teeth or mandib* or maxilla* or jaw*) and (radiograph* or radiolog* or x-ray* or xray* or imag*)).ab,ti.	54349
4	(Panoramic* or Pantomogra* or Orthopantomogra* or radiovisiograph* or Scanora* or Visualix or Visualices or Digora* or Sens-A-Ray* or Sens A Ray* or SensARay*).ab,ti.	8715
5	1 or 2 or 3 or 4	71261
6	exp trabecular bone/	17957
7	exp bone density/	80632
8	(Cancellous or spongy or trabecul* or Intertrabecul* or Bone densit* or Bone mineral densit* or BMD or Bone mineral content* or Bone texture*).ab,ti.	126297
9	6 or 7 or 8	148006
10	5 and 9	2798
11	(animal not (animal and human)).sh.	1402293
12	10 not 11	2545
13	limit 12 to yr="1990 -Current"	2444
14	limit 13 to (danish or english or finnish or norwegian or swedish)	2269
15	limit 14 to (article or conference paper or "review")	1960

Database: The Cochrane Library

Date: 2018-04-13

No. of results: 267

Cochrane Reviews (1)

Trials (266)

#	Searches	Results
#1	MeSH descriptor: [Radiography, Dental] explode all trees	458
#2	((Dental or Intraoral or Intra-oral or periapical or peri-apical or Periapex or Peri-apex or tooth or teeth or mandib* or maxilla* or jaw*) and (radiograph* or radiolog* or x-ray* or xray* or imag*)):ti,ab,kw (Word variations have been searched)	3697
#3	Panoramic* or Pantomogra* or Orthopantomogra* or radiovisiograph* or Scanora* or Visualix or Visualices or Digora* or Sens-A-Ray* or Sens A Ray* or SensARay*:ti,ab,kw (Word variations have been searched)	296
#4	#1 or #2 or #3	3782
#5	MeSH descriptor: [Cancellous Bone] explode all trees	8
#6	MeSH descriptor: [Bone Density] explode all trees	4591
#7	Cancellous or spongy or trabecul* or Intertrabecul* or "Bone densit*" or "Bone mineral densit*" or BMD or "Bone mineral content*" or "Bone texture*":ti,ab,kw (Word variations have been searched)	11008
#8	#5 or #6 or #7	11008
#9	#4 and #8 Publication Year from 1990 to 2018	267

The websites of **SBU** and **Folkhelseinstituttet** were visited 2018-04-13.
Nothing relevant to the question at issue was found.

Reference lists

A comprehensive review of reference lists brought 10 new records.

Reference lists

Included studies:

Jonasson G, Billhult A. Mandibular bone structure, bone mineral density, and clinical variables as fracture predictors: a 15-year follow-up of female patients in a dental clinic. *Oral Surg Oral Med Oral Pathol Oral Radiol.* 2013;116(3):362-8.

Sundh V, Hange D, Ahlqwist M, Hakeberg M, Lissner L, Jonasson G. FRAX and mandibular sparse trabeculation as fracture predictors: a longitudinal study from 1980 to 2002. *Eur J Oral Sci.* 2017;125(2):135-40.

Verheij JG, Geraets WG, van der Stelt PF, Horner K, Lindh C, Nicopoulou-Karayianni K, et al. Prediction of osteoporosis with dental radiographs and age. *Dentomaxillofac Radiol.* 2009;38(7):431-7.

Excluded studies:

Ay S, Gursoy UK, Erselcan T, Marakoglu I. Assessment of mandibular bone mineral density in patients with type 2 diabetes mellitus. *Dentmaxillofac Radiol.* 2005;34(6):327-31.

Engel MB, Rosenberg HM, Jordan SL, Holm K. Radiological evaluation of bone status in the jaw and in the vertebral column in a group of women. *Gerodontology.* 1994;11(2):86-92.

Geraets WG, Lindh C, Verheij H. Sparseness of the trabecular pattern on dental radiographs: visual assessment compared with semi-automated measurements. *Br J Radiol.* 2012;85(1016):e455-60.

Geraets WG, Verheij JG, van der Stelt PF, Horner K, Lindh C, Nicopoulou-Karayianni K, et al. Prediction of bone mineral density with dental radiographs. *Bone.* 2007;40(5):1217-21.

Geraets WG, Verheij JG, van der Stelt PF, Horner K, Lindh C, Nicopoulou-Karayianni K, et al. Selecting regions of interest on intraoral radiographs for the prediction of bone mineral density. *Dentmaxillofac Radiol.* 2008;37(7):375-9.

Hedstrom L, Baigi A, Bergh H. The relation between bone mineral density in the heel and pixel intensity in the mandibular jaw bone among elderly women. *Dentmaxillofac Radiol.* 2010;39(7):409-13.

Jonasson G. Bone mass and trabecular pattern in the mandible as an indicator of skeletal osteopenia: a 10-year follow-up study. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2009;108(2):284-91.

Jonasson G. Five-year alveolar bone level changes in women of varying skeletal bone mineral density and bone trabeculation. *Oral Surg Oral Med Oral Pathol Oral Radiol.* 2015;120(1):86-93.

Jonasson G, Bankvall G, Kiliaridis S. Estimation of skeletal bone mineral density by means of the trabecular pattern of the alveolar bone, its interdental thickness, and the bone mass of the mandible. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2001;92(3):346-52.

Jonasson G, Jonasson L, Kiliaridis S. Changes in the radiographic characteristics of the mandibular alveolar process in dentate women with varying bone mineral density: a 5-year prospective study. *Bone.* 2006;38(5):714-21.

Jonasson G, Jonasson L, Kiliaridis S. Skeletal bone mineral density in relation to thickness, bone mass, and structure of the mandibular alveolar process in dentate men and women. *OraEur J Oral Sci.* 2007;115(2):117-23.

Jonasson G, Sundh V, Ahlqwist M, Hakeberg M, Bjorkelund C, Lissner L. A prospective study of mandibular trabecular bone to predict fracture incidence in women: a low-cost screening tool in the dental clinic. *Bone.* 2011;49(4):873-9.

Jonasson G, Sundh V, Hakeberg M, Hassani-Nejad A, Lissner L, Ahlqwist M. Mandibular bone changes in 24 years and skeletal fracture prediction. *Clin Oral Investig.* 2013;17(2):565-72.

Kathirvelu D, Anburajan M. Prediction of low bone mass using a combinational approach of cortical and trabecular bone measures from dental panoramic radiographs. *Proc Inst Mech Eng H.* 2014;228(9):890-8.

Kavitha MS, An SY, An CH, Huh KH, Yi WJ, Heo MS, et al. Texture analysis of mandibular cortical bone on digital dental panoramic radiographs for the diagnosis of osteoporosis in Korean women. *Oral Surg Oral Med Oral Pathol Oral Radiol.* 2015;119(3):346-56.

Kavitha MS, Ganesh Kumar P, Park SY, Huh KH, Heo MS, Kurita T, et al. Automatic detection of osteoporosis based on hybrid genetic swarm fuzzy classifier approaches. *Dentmaxillofac Radiol.* 2016;45(7):20160076.

Khojastehpour L, Mogharrabi S, Dabbaghmanesh MH, Iraj Nasrabadi N. Comparison of the mandibular bone densitometry measurement between normal, osteopenic and osteoporotic postmenopausal women. *J Dent (Tehran).* 2013;10(3):203-9.

Koh KJ, Park HN, Kim KA. Prediction of age-related osteoporosis using fractal analysis on panoramic radiographs. *Imaging Sci Dent.* 2012;42(4):231-5.

Law AN, Bollen AM, Chen SK. Detecting osteoporosis using dental radiographs: a comparison of four methods. *J Am Dent Assoc.* 1996;127(12):1734-42.

Lee BD, White SC. Age and trabecular features of alveolar bone associated with osteoporosis. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2005;100(1):92-8.

- Leite AF, de Souza Figueiredo PT, Caracas H, Sindeaux R, Guimaraes ATB, Lazarte L, et al. Systematic review with hierarchical clustering analysis for the fractal dimension in assessment of skeletal bone mineral density using dental radiographs. *Oral Radiol.* 2015;31(1):1-13.
- Licks R, Licks V, Ourique F, Bittencourt HR, Fontanella V. Development of a prediction tool for low bone mass based on clinical data and periapical radiography. *Dentmaxillofac Radiol.* 2010;39(4):224-30.
- Lindh C, Horner K, Jonasson G, Olsson P, Rohlin M, Jacobs R, et al. The use of visual assessment of dental radiographs for identifying women at risk of having osteoporosis: the OSTEODENT project. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2008;106(2):285-93.
- Lopez-Lopez J, Castellanos-Cosano L, Estrugo-Devesa A, Gomez-Vaquero C, Velasco-Ortega E, Segura-Egea JJ. Radiolucent periapical lesions and bone mineral density in post-menopausal women. *Gerodontology.* 2015;32(3):195-201.
- Lundstrom A, Jendle J, Stenstrom B, Toss G, Ravald N. Periodontal conditions in 70-year-old women with osteoporosis. *Swed Dent J.* 2001;25(3):89-96.
- Lurie A, Tosoni GM, Tsimikas J, Fitz Jr W. Recursive hierarchic segmentation analysis of bone mineral density changes on digital panoramic images. *Oral Surg Oral Med Oral Pathol Oral Radiol.* 2012;113(4):549-58.
- Mohajery M, Brooks SL. Oral radiographs in the detection of early signs of osteoporosis. *Oral Surg Oral Med Oral Pathol.* 1992;73(1):112-7.
- Mohammad AR, Alder M, McNally MA. A pilot study of panoramic film density at selected sites in the mandible to predict osteoporosis. *Int J Prosthodont.* 1996;9(3):290-4.
- Nackaerts O, Jacobs R, Devlin H, Pavitt S, Bleyen E, Yan B, et al. Osteoporosis detection using intraoral densitometry. *Dentmaxillofac Radiol.* 2008;37(5):282-7.
- Roberts MG, Graham J, Devlin H. Image texture in dental panoramic radiographs as a potential biomarker of osteoporosis. *IEEE Trans Biomed Eng.* 2013;60(9):2384-92.
- Scheibel PC, Ramos AL, Iwaki LC. Is there correlation between alveolar and systemic bone density? *Dental Press J Orthod.* 2013;18(5):78-83.
- Sindeaux R, Figueiredo PT, de Melo NS, Guimaraes AT, Lazarte L, Pereira FB, et al. Fractal dimension and mandibular cortical width in normal and osteoporotic men and women. *Maturitas.* 2014;77(2):142-8.
- Takaishi Y, Arita S, Honda M, Sugishita T, Kamada A, Ikeo T, et al. Assessment of alveolar bone mineral density as a predictor of lumbar fracture probability. *Adv Ther.* 2013;30(5):487-502.
- Tosoni GM, Lurie AG, Cowan AE, Burlison JA. Pixel intensity and fractal analyses: detecting osteoporosis in perimenopausal and postmenopausal women by using digital panoramic images. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2006;102(2):235-41.
- White SC, Atchison KA, Gornbein JA, Nattiv A, Paganini-Hill A, Service SK, et al. Change in mandibular trabecular pattern and hip fracture rate in elderly women. *Dentmaxillofac Radiol.* 2005a;34(3):168-74.
- White SC, Rudolph DJ. Alterations of the trabecular pattern of the jaws in patients with osteoporosis. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 1999;88(5):628-35.

White SC, Taguchi A, Kao D, Wu S, Service SK, Yoon D, et al. Clinical and panoramic predictors of femur bone mineral density. *Osteoporos Int.* 2005b;16(3):339-46.

Yasar F, Akgunlu F. The differences in panoramic mandibular indices and fractal dimension between patients with and without spinal osteoporosis. *Dentmaxillofac Radiol.* 2006;35(1):1-9.

Yilmaz HH, Akgunlu F, Aydin U. Comparison of trabecular pattern in healthy and osteoporotic women. *Gulhane Med J.* 2004;46(3):200-4.

Other references:

Andersson, Pål (2007). Strålmiljön i Sverige. SSI-rapport, 0282-4434 ; 2007:02. Stockholm: Statens strålskyddsinstitut. Libris 10665865. Accessed 16 Jan 2019.

Atkins D, Best D, Briss PA, Eccles M, Falck-Ytter Y, Flottorp S, et al. GRADE Working Group. Grading quality of evidence and strength of recommendations. *BMJ.* 2004 Jun 19;328(7454):1490-4.

Blake GM, Naeem M, Boutros M. Comparison of effective dose to children and adults from dual X-ray absorptiometry examinations. *Bone.* 2006;38(6):935-42.

Bolland MJ, Siu AT, Mason BH, Horne AM, Ames RW, Grey AB, et al. Evaluation of the FRAX and Garvan fracture risk calculators in older women. *J Bone Miner Res.* 2011;26(2):420-7.

Borgström F, Zethraeus N, Johnell O, Lidgren L, Ponzer S, Svensson O, et al. Costs and quality of life associated with osteoporosis-related fractures in Sweden. *Osteoporos Int.* 2006;17(5):637-50.

Brasileiro CB, Chalub LLFH, Abreu MHNG, Barreiros ID, Amaral TMP, Kakehasi AM, et al. Use of cone beam computed tomography in identifying postmenopausal women with osteoporosis. *Arch Osteoporos.* 2017;12(1):26.

Ensrud KE, Lui LY, Taylor BC, Schousboe JT, Donaldson MG, Fink HA, et al. A comparison of prediction models for fractures in older women: is more better? *Arch Intern Med.* 2009;169(22):2087-94.

Fraser LA, Langsetmo L, Berger C, Ioannidis G, Goltzman D, Adachi JD, et al. Fracture prediction and calibration of a Canadian FRAX® tool: a population-based report from CaMos. *Osteoporos Int.* 2011;22(3):829-37.

FRAX® Fracture Risk Assessment Tool [Internet]. Sheffield: University of Sheffield, Centre for Metabolic Bone Diseases; 2008 - [cited 2018 Oct 30]. Available from: <https://www.sheffield.ac.uk/FRAX/tool.aspx?country=5>

Geraets WG, Lindh C, Verheij H. Sparseness of the trabecular pattern on dental radiographs: visual assessment compared with semi-automated measurements. *Br J Radiol.* 2012;85(1016):e455-60.

Geraets WG, Verheij JG, van der Stelt PF, Horner K, Lindh C, Nicopoulou-Karayianni K, et al. Prediction of bone mineral density with dental radiographs. *Bone.* 2007;40(5):1217-21.

GRADE Working Group. [Internet]. [Place unknown]: GRADE Working Group, c2004-2018 [cited 2018 Oct 31]. Available from: <http://www.gradeworkinggroup.org>

Huja SS, Fernandez SA, Hill KJ, Li Y. Remodeling dynamics in the alveolar process in skeletally mature dogs. *Anat Rec A Discov Mol Cell Evol Biol.* 2006;288(12):1243-9.

Jonasson G, Sundh V, Ahlqwist M, Hakeberg M, Björkelund C, Lissner L. A prospective study of mandibular trabecular bone to predict fracture incidence in women: a low-cost screening tool in the dental clinic. *Bone*. 2011;49(4):873-9.

Kanis JA. *Osteoporosis*. Oxford: Blackwell Science; 1994.

Kanis JA, Oden A, Johnell O, Johansson H, De Laet C, Brown J, et al. The use of clinical risk factors enhances the performance of BMD in the prediction of hip and osteoporotic fractures in men and women. *Osteoporos Int*. 2007;18(8):1033-46.

Kanis JA, Oden A, Johnell O, Jonsson B, de Laet C, Dawson A. The burden of osteoporotic fractures: a method for setting intervention thresholds. *Osteoporos Int*. 2001;12(5):417-27.

Kathirvelu D, Anburajan M. Prediction of low bone mass using a combinational approach of cortical and trabecular bone measures from dental panoramic radiographs. *Proc Inst Mech Eng H*. 2014;228(9):890-8.

Kavitha MS, An SY, An CH, Huh KH, Yi WJ, Heo MS, et al. Texture analysis of mandibular cortical bone on digital dental panoramic radiographs for the diagnosis of osteoporosis in Korean women. *Oral Surg Oral Med Oral Pathol Oral Radiol*. 2015;119(3):346-56.

Kavitha MS, Ganesh Kumar P, Park SY, Huh KH, Heo MS, Kurita T, et al. Automatic detection of osteoporosis based on hybrid genetic swarm fuzzy classifier approaches. *Dentomaxillofac Radiol*. 2016;45(7):20160076.

Koh KJ, Park HN, Kim KA. Prediction of age-related osteoporosis using fractal analysis on panoramic radiographs. *Imaging Sci Dent*. 2012;42(4):231-5.

Lee BD, White SC. Age and trabecular features of alveolar bone associated with osteoporosis. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 2005;100(1):92-8.

Leslie WD, Lix LM, Johansson H, Oden A, McCloskey E, Kanis JA; et al. Independent clinical validation of a Canadian FRAX tool: fracture prediction and model calibration. *J Bone Miner Res*. 2010;25(11):2350-8.

Licks R, Licks V, Ourique F, Radke Bittencourt H, Fontanella V. Development of a prediction tool for low bone mass based on clinical data and periapical radiography. *Dentomaxillofac Radiol*. 2010;39(4):224-30.

Lindh C, Horner K, Jonasson G, Olsson P, Rohlin M, Jacobs R, et al. The use of visual assessment of dental radiographs for identifying women at risk of having osteoporosis: the OSTEODENT project. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 2008;106(2):285-93.

Ludlow JB, Davies-Ludlow LE, White SC. Patient risk related to common dental radiographic examinations: the impact of 2007 International Commission on Radiological Protection recommendations regarding dose calculation. *J Am Dent Assoc*. 2008;139(9):1237-43.

Lurie A, Tosoni GM, Tsimikas J, Walker F Jr. Recursive hierarchic segmentation analysis of bone mineral density changes on digital panoramic images. *Oral Surg Oral Med Oral Pathol Oral Radiol*. 2012;113(4):549-58.e1.

Moher D, Liberati A, Tetzlaff J, Altman DG; PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med*. 2009 Jul 21;6(7):e1000097.

Mostafa RA, Arnout EA, Abo El-Fotouh MM. Feasibility of cone beam computed tomography radiomorphometric analysis and fractal dimension in assessment of postmenopausal osteoporosis in correlation with dual X-ray absorptiometry. *Dentomaxillofac Radiol.* 2016;45(7):20160212.

National Clinical Guideline Centre (UK). 4. Risk assessment tools (FRAX, QFracture, BMD). In: *Osteoporosis: fragility fracture risk: osteoporosis: assessing the risk of fragility fracture.* (NICE Clinical Guidelines, No. 146.). London: Royal College of Physicians (UK); 2012. [cited 2018 Dec 17]. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK115321/>

Pasco JA, Seeman E, Henry MJ, Merriman EN, Nicholson GC, Kotowicz MA. The population burden of fractures originates in women with osteopenia, not osteoporosis. *Osteoporos Int.* 2006;17(9):1404-9.

QUADAS-2: a revised tool for the quality assessment of diagnostic accuracy studies. [Internet]. [cited 2018 Jun 13] Available from: <https://www2.sahlgrenska.se/upload/SU/HTA-centrum/Hj%C3%A4lpmedel%20under%20projektet/QUADAS%202.pdf>

Sahlgrenska Universitetssjukhuset. Osteoporosmottagning [Internet]. Göteborg: Sahlgrenska Universitetssjukhuset; [updated 2018 Aug 30; cited 2018 Nov 20]. Available from: <http://www.sahlgrenska.se/omrade-3/geriatrik/enheter/osteoporosmottagning/>

Sanders KM, Nicholson GC, Watts JJ, Pasco JA, Henry MJ, Kotowicz MA, et al. Half the burden of fragility fractures in the community occur in women without osteoporosis. When is fracture prevention cost-effective? *Bone.* 2006;38(5):694-700.

Socialstyrelsen. Nationella riktlinjer för rörelseorganens sjukdomar 2012: osteoporos, artros, inflammatorisk ryggskjutdom och ankyloserande spondylit, psoriasisartrit och reumatoid artrit : stöd för styrning och ledning. Stockholm: Socialstyrelsen; 2012. [cited 2018 Nov 20]. Available from: <http://www.socialstyrelsen.se/Lists/Artikelkatalog/Attachments/18665/2012-5-1.pdf>

Socialstyrelsen. Vård vid rörelseorganens sjukdomar: rekommendationer, bedömningar och sammanfattning. Stockholm: Socialstyrelsen; 2014.

Tu KN, Lie JD, Wan CKV, Cameron M, Austel AG, Nguyen JK, et al. Osteoporosis: A Review of Treatment Options. *P T.* 2018;43(2):92-104.

White SC, Atchison KA, Gornbein JA, Nattiv A, Paganini-Hill A, Service SK, et al. Change in mandibular trabecular pattern and hip fracture rate in elderly women. *Dentomaxillofac Radiol.* 2005a;34(3):168-74.

Whiting PF, Rutjes AW, Westwood ME, Mallett S, Deeks JJ, Reitsma JB, et al.; QUADAS-2 Group. QUADAS-2: a revised tool for the quality assessment of diagnostic accuracy studies. *Ann Intern Med.* 2011;155(8):529-36.

Västra Götalandsregionen. Osteoporosrelaterade frakturer (Dnr RS 454-2013). [Internet]. [Place unknown]: Västra Götalandsregionen; 2013 [cited 2018 Oct 30]. Available from: [https://alfresco.vgregion.se/alfresco/service/vgr/storage/node/content/workspace/SpacesStore/c2183b19-283d-4d61-a77f-37bdf08dafa0/Osteoporosrelaterade%20frakturer%20\(v%C3%A5rdprocess\).pdf?a=false&guest=true](https://alfresco.vgregion.se/alfresco/service/vgr/storage/node/content/workspace/SpacesStore/c2183b19-283d-4d61-a77f-37bdf08dafa0/Osteoporosrelaterade%20frakturer%20(v%C3%A5rdprocess).pdf?a=false&guest=true)

Västra Götalandsregionen. Regional medicinsk riktlinje – läkemedel. Osteoporos [Internet]. Västra Götalandsregionen; 2018. [cited 2019 Jan 10]. Available from: <https://alfresco.vgregion.se/alfresco/service/vgr/storage/node/content/3351/Osteoporos.pdf?a=false&guest=true>

Västra Götalandsregionen, Vårddatabas Vega, 2019.

Appendix 2 – Characteristics of included studies

Project: Trabecular bone pattern assessment in dental radiographs for screening of fracture risk

Author Year Country	Study design	Study duration (years)	Study groups; Index vs reference	Patients (n)	Mean age (years)	Men (%)	Outcome variables
Jonasson & Billhult 2013, Sweden.	Cohort	15 years, 1996-2011	Intraoral radiograph vs. FRAX and/or BMD (by DXA)	136	64.1 SD 11.2 Range 35-94	0	Fracture risk
Sundh et al 2017, Sweden.	Cohort	10 years 1980-1992 10 years 1992-2002	Panoramic radiograph vs. FRAX	499 (1980-1992) 412 (1992-2002)	Range 50-66 (1980-1992) Range 62-78 (1992-2002)	0	Fracture risk
Verheij et al 2009, Belgium, Greece, The Netherlands, Sweden, United Kingdom	Cross sectional	Not reported	Panoramic and intraoral radiographs vs. BMD (by DXA)	607	54.8 SD 6.1 Range 45-71	0	Sensitivity and specificity for osteoporosis measured with DXA*

BMD: Bone mineral density, DXA: Dual-energy X-ray absorptiometry, FRAX: Fracture risk assessment tool* i.e. DXA BMD T-score ≤ -2.5

Appendix 3.

Project: Trabecular bone pattern assessment in dental radiographs for screening of fracture risk

Excluded articles

Author, year	Reason for exclusion
Ay, S., 2005	Wrong I
Engel, M.B., 1994	Wrong I
Geraets W.G., 2012	Wrong O
Geraets W.G., 2007	Wrong O
Geraets W.G., 2008	Wrong O
Hedström L., 2010	Wrong C
Jonasson G., 2009	Wrong O
Jonasson G., 2015	Wrong O
Jonasson G., 2001	Wrong O
Jonasson G., 2006	Wrong O
Jonasson G., 2007	Wrong O
Jonasson G., 2011	Duplicate material of Jonasson G., 2013
Jonasson G., 2013	Case series not reporting risks or complications
Kathirvelu D and Anburajan M., 2014	Wrong O
Kavitha M.S., 2015	Wrong I
Kavitha M.S., 2016	Wrong I
Khojastehpour L., 2013	Wrong I
Koh K.J., 2012	Wrong P
Law A.N., 1996	Wrong P
Lee B.D., 2005	Data not extractable
Leite A.F., 2015	Wrong I
Licks R., 2010	Wrong O
Lindh C., 2008	Wrong P
Lopez- Lopez, J., 2015	Wrong P
Lundstrom A., 2001	Wrong P
Lurie A., 2012	Wrong I
Mohajery M., 1992	Wrong I
Mohammad A.R., 1996	Wrong P
Nackaerts, O., 2008	Wrong P
Roberts M.G., 2013	Wrong I
Scheibel, P.C., 2013	Wrong I
Sindeaux R., 2014	Wrong P
Takishi Y., 2013	Wrong I
Tosoni G.M., 2006	Wrong P
White S.C., 1999	Wrong P
White S.C., 2005a	Case series not reporting risks or complications
White S.C., 2005b	Wrong P
Yasar F., 2006	Wrong O
Yilmaz H.H., 2004	Wrong P

Appendix 4.1

Project: Trabecular bone pattern assessment in dental radiographs for screening of fracture risk

Outcome variable: Fracture prediction

* + No or minor problems
 ? Some problems
 - Major problems

Author year country	Study design	Number of patients n=	Withdrawals - dropouts	Results		Comments	Directness *	Study limitations *	Precision *
				Index test (Trabecular bone pattern assessment in dental radiographs)	Reference test (FRAX, BMD)				
Jonasson & Billhult 2013, Sweden	Observational, prospective cohort study	1996 n=166 2011 n=136 replied (14 died)	n=30	<p><i>Fracture prediction:</i></p> <p><u>Visual evaluation, sparse trabeculation</u> RR 1.52 95% CI: 0.56 to 4.11, n.s. AR 35%</p> <p><u>Jaw-X (software)</u> RR 1.87 95% CI: 0.73 to 4.71, n.s.</p> <p><u>Bone texture (software)</u> RR 4.74 95% CI: 1.49 to 15.04, p<0.05</p>	<p><i>Fracture prediction:</i></p> <p><u>FRAX with BMD²</u> RR 2.53 95% CI: 1.13 to 5.63, p<0.05</p> <p><u>FRAX without BMD</u> RR 1.34 95% CI: 0.56 to 3.20, n.s.</p> <p><u>BMD</u> RR 2.37 95% CI: 0.93 to 6.02, n.s.</p>	<p>BMD: DXA forearm</p> <p>To calculate the RR, the FRAX value was dichotomised, where $\geq 7\%$ indicated increased risk for fracture.</p> <p>Blinded evaluation (according to author Jonasson). Data included in FRAX, from 1996, were analysed retrospectively in 2011 (alcohol consumption was not included in FRAX).</p> <p>6 women had T-score ≤ -2.5</p>	+	?	-
Sundh <i>et al.</i> , 2017 Sweden	Observational, prospective cohort study	1980/ 1981 n=499 1992/ 1993 n=412	No dropouts	<p><i>Fracture prediction 1980-1992:</i></p> <p><u>Visual evaluation, sparse trabeculation</u> RR 2.09 95% CI: 1.3 to 3.5</p> <p>AR: 23.0%</p> <p>OR: 3.03 95% CI: 1.81 to 5.08</p>	<p><i>Fracture prediction 1980-1992:</i></p> <p><u>FRAX >15%</u> RR 1.85 95% CI: 0.7 to 5.6</p> <p>AR: 36.7%</p> <p>OR: 4.35 95% CI: 2.28 to 8.31</p>	<p>Blinded, according to author. Information on fractures in parents was not available and not included in the FRAX calculation.</p> <p>397 participated in both (1980-1993 and 1992-2002) examinations.</p>	+	+/?	-

Appendix 4.1

Project: Trabecular bone pattern assessment in dental radiographs for screening of fracture risk

Outcome variable: Fracture prediction

* + No or minor problems
 ? Some problems
 - Major problems

Author year country	Study design	Number of patients n=	Withdrawals - dropouts	Results		Comments	Directness *	Study limitations *	Precision *
				Index test (Trabecular bone pattern assessment in dental radiographs)	Reference test (FRAX, BMD)				
				<i>Fracture prediction 1992-2002:</i> <u>Visual evaluation, sparse trabeculation</u> RR 3.7 95% CI: 2.2 to 6.4 AR: 26.1% OR: 4.57 95% CI: 2.47 to 8.45	<i>Fracture prediction 1992-2002:</i> <u>FRAX >15%</u> RR 4.1 95% CI: 2.4 to 7.2 AR: 26.9% OR: 5.28 95% CI: 2.84 to 9.94				

AR = Absolute risk, BMD = bone mineral density, Bone texture: a computer software, CI = confidence interval, FRAX = fracture risk assessment tool, Jaw-X: a computer software, based on an image processing algorithm, OR = Odds ratio, RR = relative risk.

Appendix 4.2

Project: Trabecular bone pattern assessment in dental radiographs for screening of fracture risk

Outcome variable: Sensitivity and specificity for osteoporosis measured with DXA*

* + No or minor problems
 ? Some problems
 - Major problems

Author year country	Study design	Number of patients n=	Withdrawals - dropouts	Results		Comments	Directness *	Study limitations *	Precision *
				Index test	Reference test				
Verheij <i>et al.</i> , 2009 Five countries in Europe: The Netherlands, Belgium, Greece, UK, Sweden.	Observational cross-sectional study	n=671	n=64	<i>Morphological digital analysis of trabecular bone pattern (software)</i> <u>Trabecular pattern</u> For reference test: Sensitivity 0.70 (95% CI 0.62 to 0.78) Specificity 0.69 (95% CI 0.64 to 0.73)	<i>DXA hip, spine</i> <u>BMD</u> T-score \leq -2.5	Convenience sample. No pre-determined threshold. Split of Index variable to obtain maximal sensibility and specificity, i.e. no blinding. Mean age 54.8 years (SD 6.1) Regions of interest on the radiographs were chosen by preference. No indication of blinding.	+	?	-

BMD = Bone mineral density, CI = Confidence interval, DXA = Dual-energy X-ray absorptiometry.

* i.e. DXA BMD T-score \leq -2.5

Region Västra Götaland, HTA-centrum

Health Technology Assessment
Regional activity-based HTA



HTA

Health technology assessment (HTA) is the systematic evaluation of properties, effects, and/or impacts of health care technologies, i.e. interventions that may be used to promote health, to prevent, diagnose or treat disease or for rehabilitation or long-term care. It may address the direct, intended consequences of technologies as well as their indirect, unintended consequences. Its main purpose is to inform technology-related policymaking in health care.

To evaluate the quality of evidence the Centre of Health Technology Assessment in Region Västra Götaland is currently using the GRADE system, which has been developed by a widely representative group of international guideline developers. According to GRADE the level of evidence is graded in four categories:

High quality of evidence	= (GRADE ⊕⊕⊕⊕)
Moderate quality of evidence	= (GRADE ⊕⊕⊕⊖)
Low quality of evidence	= (GRADE ⊕⊕⊖⊖)
Very low quality of evidence	= (GRADE ⊕⊖⊖⊖)

In GRADE there is also a system to rate the strength of recommendation of a technology as either “strong” or “weak”. This is presently not used by the Centre of Health Technology Assessment in Region Västra Götaland. However, the assessments still offer some guidance to decision makers in the health care system. If the level of evidence of a positive effect of a technology is of high or moderate quality it most probably qualifies to be used in routine medical care. If the level of evidence is of low quality the use of the technology may be motivated provided there is an acceptable balance between benefits and risks, cost-effectiveness and ethical considerations. Promising technologies, but a very low quality of evidence, motivate further research but should not be used in everyday routine clinical work.

Christina Bergh, Professor, MD.
Head of HTA-centrum

